

Species Composition and Biomass Production in Two Communal Coastal Rangelands of the Eastern Cape Province, South Africa

Matshawule, S.

Department of Rural Development & Agrarian Reform, Chris Hani district, P.O. Box 112, Queenstown, 5320, Eastern Cape, Republic of South Africa.
Email: matshawule.sinethemba@yahoo.com



Abstract

This study investigated species composition and biomass production of herbaceous plants in two communal rangelands surrounding homesteads areas. In each study area, the rangelands were divided based on the distance from homesteads into near (up to 1 km), middle (> 1–2 km) and far (> 2–3 km) sites. On each site a HVU of 50 m x 20 m was used and six 0.25 m² quadrants were laid randomly on each HVU. Herbaceous species found within a quadrant were identified, counted and their height and tuft were measured. Herbaceous species were also harvested, bulked and placed into well labelled brown paper bags and oven-dried for 48 hours at 60 °C to determine biomass production. Herbaceous species were also classified according to their palatability ecological status and life form. Data were collected for two seasons over 2014/15 (winter and summer). A total of 20 herbaceous species were identified in the study areas. Of these 17 were grass species. The most common or dominant grass species were *T. triandra*, *C. dactylon*, *E. capensis*, *E. plana* and *S. africanus*. At Dyamdyam *T. triandra* showed the greatest frequency of occurrence at far site than middle and near homesteads sites. The frequency occurrence of *S. africanus* at Machibi was relatively similar in all the study sites. Density of herbaceous species was significantly lowest on near and bottom sites at Dyamdyam and Machibi, respectively. In both winter and summer, grazing site far and top from homestead had greatest ($P < 0.05$) biomass production at Dyamdyam and Machibi respectively. The biomass production was significantly highest in the summer than the winter across the study sites in both study areas. It can be concluded that composition of herbaceous species and biomass production are significantly dependent on distance and topography from homesteads and seasons. Therefore, any rangeland management practices in communal grazing lands should consider these factors in to consideration during the planning of development progress.

Keywords

Distance Gradient, Livestock, Topography, Summer, Soil, Vegetation, Winter

<http://abr.damray.com>

OPEN ACCESS

Received: September 25, 2019

Accepted: October 29, 2019

Published: April 14, 2020

Copyright: ©2020 Matshawule, S. et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

1. Introduction

Communal rangelands are used mainly as sources of forage for livestock and collection of wood for fire and building (Everson & Hatch, 1999). These rangelands have different species composition and biomass productions

(Oztas et al., 2003). Species composition and biomass production on communal rangeland vary according to grazing system (Shackleton, 1993), seasonal variation (Angasa & Oba, 2010), soil type (O' Farrell et al., 2007), topography (Lesoli, 2008) and grazing intensity (Smet & Ward, 2003; Maki et al., 2007).

Species composition is defined as the relative frequency of occurrence of heterogeneous herbaceous species in a rangeland (Trollope et al., 1990). In addition, it is one of the factors that indicate the rangeland condition because herbaceous species differ significantly in their acceptability, ecological status, life form and response to grazing (Abule et al., 2007). It has been reported that, high grazing pressure on natural rangelands causes changes in species composition (Maki et al., 2007). Decreaser species disappear in response to heavy grazing and are replaced by increaser and invader species that are less palatable and adapted to withstand over utilization (Sisay & Baars, 2002). Most rangelands in communal areas of South Africa are continuously grazed due to the absence of fence and this affects distribution of herbaceous species (Shackleton, 1993). In addition, high grazing intensity can alter vegetation from being dominated by perennial species to being dominated by annual species due to the high stocking rate on grazing lands (Smet & Ward, 2003). Other researchers at species composition change is determined more by rainfall rather than by grazing pressure. In summer there is more rainfall that promotes height, diameter and basal cover of grass species.

Biomass production is a total dry matter of living material which is actively and structurally functional in a given area and is normally used as source of energy for livestock and for fuel (Bond-Lamberty et al., 2002). According to Angasa & Oba (2010), biomass production during the dry season is lower than during the wet season. This is allied with high amount of rainfall and availability of moisture during wet season than dry season (Angasa & Oba, 2010). The part of rangelands that is heavily grazed for long period of time has lower biomass production (Shackleton, 1993). In addition, biomass production in silt and clay soil is higher than in sandy soil due to the high organic matter and minerals in clay and silt soils than sandy soils (O' Farrell et al., 2007).

Landscape positions have significant effects on the biomass production with several reports showing bottomlands having higher biomass production than slope and top lands (Coronato & Bertiller, 1996; Lesoli, 2008). In contrast, other researchers reported bottomlands have lower biomass due to the presence of heavy grazing pressure because of easily access by livestock than the plants found on top lands (Senft et al., 1985; Belsky & Blumenthal, 1997; Lesoli, 2008). Studies that are conducted on communal coastal rangelands of South Africa to investigate the change in species composition and biomass production along the gradient distance from homestead are limited. There is also no information regarding the seasonal variations in the above variables along distance gradient from homesteads. The availability of such information would contribute towards of planning management and rehabilitation programs in the communal rangelands to improve fodder production for livestock. Therefore, the objectives of this study were 1) to assess herbaceous species distribution and biomass production along distance gradient from homesteads in two communal rangelands, 2) to investigate the effects of season on biomass production and species distribution.

2. Methods and Materials

2.1. Selection of Study Sites and Layout

Two long transect radiating away from the homesteads in each communal area was established and divided into three main grazing sites. The transects were established in the directions of the main grazing activities. Dyamdyam is situated on relatively plain land and; hence the rangeland was simply divided according to the distance from homesteads into near (up to 1 km), middle (> 1–2 km) and far (> 2–3 km) sites. Machibi is set on a gentle steep slope, and therefore three grazing sites were identified to represent the bottom (near), slope (middle) and top (far) sites from homestead. On each study site a HVU of 20 m x 50 m was marked to record vegetation data from six 0.25 m² quadrants that were laid randomly.

2.2. Data Collection

Data on herbaceous species composition, tuft diameter, height, density and biomass were collected from two different directions where the livestock are grazing in each study area. In each 0.25 m² quadrants herbaceous species found within a quadrant were identified, counted and recorded. Height and tuft diameter of herbaceous species were measured using 30 cm ruler and recorded. Total density was obtained by adding all the herbaceous

species found in each site. Density for each herbaceous species found in study areas were then calculated as the percentage to obtain the frequency of occurrence of each herbaceous species. Herbaceous species found within a quadrant were harvested, bulked and placed into well label brown paper bags. Harvested herbaceous samples were then oven-dried for 24 hours at 60 °C. Dried grass samples were weighed to determine dry matter (DM) production. Data was collected in winter and summer in 2014 and 2015.

2.3. Species Classification

Classification of herbaceous species was based on the succession theory described by Dyksterhuis (1994) and on the ecological information for the arid to semi-arid region of South Africa (Vorster, 1982). Herbaceous species were classified according to their ecological status and response to grazing such as: (i) highly palatable: those species occur in rangeland that is in good condition and decrease with heavy grazing (decreaser species); (ii) palatable species: those species occur in rangeland in good and increase with moderate overgrazing (increaser IIa); (iii) less palatable species: those species which occur in rangeland in good condition and increase with high overgrazing (increaser IIb and IIc) and (iv) poorly palatable: those species which occur in rangeland in poor condition and increase with extreme overgrazing (invaders).

Herbaceous species were also grouped according to their life form as annual and perennial (Van Oudtshoorn, 1992). Grasses were further identified to species level, while other herbaceous plants belonging to other families were categorised as forbs, sedges and Karoo species.

2.4. Statistical Analyses

Data on biomass and species composition were analysed using General Linear Model (GLM) procedure of SAS (2010). Mean separation was done using PDIF option of SAS (2010). Data analyses for two study areas were done separately because they had different landscape, geology, altitude and vegetation types. Descriptive statistics such as percentage and means were used where applicable. Species composition, biomass, density and height of herbaceous species show significant interaction between site and seasons in both study areas.

3. Results

3.1. Ecological Status, Life Form, Palatability and Frequency of Occurrence of Herbaceous Species

A total of 20 herbaceous species were identified in the study areas. Three of the identified herbaceous species were categorized as forb, sedge, and Karoo species whereas 17 were grasses. In terms of their life form, all the identified herbaceous species were perennial (Table 1). In terms of their palatability, 6 herbaceous species were highly palatable, 1 was moderately palatable and 13 were low palatable. According to their ecological index, 4 grass species were classified as Decreaser species, 4 Increase I, 9 Increase II, 2 Increase III and 1 invader species (Table 1). Of the total grass species identified, 5 grass species were classified as most common or dominant grass species over the study sites (Table 1). These include *T. triandra*, *C. dactylon*, *E. capensis*, *E. plana* and *S. africanus*. One of the common or dominant species was highly desired by livestock (*T. triandra*).

Table 1: Ecological status, life form, palatability and frequency occurrence of herbaceous species identified in the study areas.

Grass species	ES	LF	Pa	Dyamdyam			Machibi		
				Near	Middle	Far	Bottom	Slope	Top
<i>Themeda triandra</i>	De	P	HP	LC	LC	C	C	D	D
<i>Cynodon dactylon</i>	Inc II	P	HP	LC	D	C	C	LC	C
<i>Eragrostis plana</i>	Inc II	P	LP	LC	C	C	LC	LC	D
<i>Sporobolus africanus</i>	IncII	P	LP	LC	C	D	C	C	C
<i>Tristachya leucothrix</i>	Inc I	P	LP	LC	LC	LC			
<i>Elulia vilosa</i>	Inc I	P	LP	R	LC	LC			
<i>Hyperrhenia hirta</i>	IncI	P	LP	R	+	+			

<i>Digitaria eriantha</i>	De	P	HP				R	R	D
<i>Paspalum dilatatum</i>	Inv	P	LP	LC	R	R	R	R	R
<i>Setaria megaphylla</i>	De	P	HP	LC	R	R	R	R	LC
<i>Brachiaria serrate</i>	De	P	HP	LC	R	R	R	R	R
<i>Cymbopogon excavatus</i>	IncI	P	LP	LC	R	R	R	R	R
<i>Sedges</i>	IncIII	P	LP	R	R	R			
<i>Forbs</i>	IncII	P	LP	R	LC	R	R	R	LC
<i>Karoo</i>	IncII	P	LP	R	R	R	LC	R	R

ES= ecological status, LF= life form, Pa= palatability, De= decreaser, IncI= increaser I, IncII= increaser II, Inc III= increaser III, Inv= invader, P= perennial, HP= high palatable, MP= moderate palatable, LP= low palatable, D= dominant (>15 %), C= common (10 %-15 %), LC=less common (5 %-10 %), R= rear (1 %-5 %) += present (<1 %) and - = absence

3.2. Common or dominant species

Common or dominant grass species in this study are defined as those species recorded along the distance gradient from homesteads in each study area and had >15 % (dominant) and >10 %- 15 % (common) frequency occurrence at least in one of the study sites. Results for frequency of occurrence of common or dominant grass species identified at Dyamdyam rangeland are presented in Figure 1. At Dyamdyam frequency occurrence of *T. triandra* was significantly higher on far site than middle and near sites. Along the distance gradient from homestead *C. dactylon* had the greatest occurrence on middle site followed by the far and near sites. The frequency occurrence *E. plana* was relatively similar on middle and far sites however were higher than near site. The proportion occurrence of *S. africanus* was significantly higher on far site followed by the middle and near sites.

Results for frequency occurrence of common or dominant grass species identified at Machibi rangeland are presented in Figure 2. At Machibi the frequency occurrence of *T. triandra* was significantly highest on slope site followed by the top and bottom sites. Along the landscape from homestead *E. plana* had greatest proportion occurrence on top site than slope and bottom sites. On bottom and top sites, the frequency occurrence of *C. dactylon* was greatest and lowest on slope site. Frequency occurrence of *S. africanus* has similar values on slope and top sites but was higher than bottom site.




Figure 1: Species composition (%) of common or dominant grasses based on frequency of occurrence at Dyamdyam. N= near, M= middle and F= far




Figure: 2: Species composition (%) of common or dominant grasses based on frequency of occurrence at Machibi. B= bottom, S= slope and T= top

3.3. Biomass production of herbaceous species

In winter, far grazing site from homesteads at Dyamdyam had the greatest ($P < 0.05$) biomass production followed by middle and near sites. In summer, far and middle grazing sites had higher biomass production than near site (Table 2). In winter, top grazing site at Machibi had highest ($P < 0.05$) biomass production followed by midslope and bottom sites. In summer, top and midslope grazing sites had higher biomass production than bottom site. Biomass production in all the study sites was significantly different between winter and summer season except for far and top sites at Dyamdyam and Machibi respectively (Table 2).

Table 2: Mean biomass production (kg ha^{-1}) of herbaceous species harvested from study areas.

Areas/Sites	Biomass	Biomass
<i>Dyamdyam</i>	Winter	Summer
Near	715.5 ^{Cb}	1015.0 ^{Ba}
Middle	1042.1 ^{Bb}	1127.5 ^{Aa}
Far	1106.3 ^{Aa}	1113.0 ^{Aa}
SE	67.5	67.5
<i>Machibi</i>		
Bottom	405.9 ^{Cb}	669.6 ^{Ba}
Slope	780.4 ^{Bb}	857.8 ^{Aa}
Top	809.1 ^{Aa}	840.4 ^{Aa}
SE	79.6	79.6

Lowercase superscripts are used to compare means between seasons within each site. ^{ABC}Uppercase superscripts are used to compare site averages within each season. Means with different superscript within the columns and rows are significantly different ($P < 0.05$).

3.4. Height and tuft diameter of common or dominant species

The results for plant height and tuft diameter of common or dominant grass species are presented in Table 3 and 4 respectively. The plant height and tuft diameter were significantly different across the study sites in some

common or dominant grass species, while in other species they were not significantly different between the study sites. *Themeda triandra* at Dyamdyam had the highest ($P < 0.05$) height in middle site followed by the far and near sites. Highest plant height for *C. dactylon* and *E. plana* at Dyamdyam was recorded in far site. Plant height of *S. africanus* was highest ($P < 0.05$) in middle site followed by the near and far sites.

At Machibi plant height of *T. triandra* was highest ($P < 0.05$) in the slope and top sites than the bottom site. Plant height of *E. plana* was significantly highest in slope site followed by bottom and top sites. Plant height of *S. africanus* was significantly highest in top site followed by slope and bottom sites. Plant height of *C. dactylon* was highest ($P < 0.05$) on samples collected from top site followed by bottom and slope sites.

Table 3: Mean height (cm) of common grass species found in study areas

Grass species	Dyamdy am				Machibi			
	Near	Middle	Far	SE	Bottom	Slope	Top	SE
<i>T. triandra</i>	11 ^c	18 ^a	12 ^b	2.3	9.0 ^b	11 ^a	11 ^a	2.0
<i>C. dactylon</i>	13 ^c	15 ^b	17 ^a	2.3	12 ^b	9.3 ^c	14 ^a	4.0
<i>E. plana</i>	14 ^c	15 ^b	20 ^a	2.3	11 ^b	13 ^a	9.0 ^c	2.1
<i>S. africanus</i>	13 ^b	20 ^a	11 ^c	2.3	9.0 ^b	9.2 ^b	11 ^a	1.9

^{abc}Different superscripts for each species in a row denote significant differences at $p < 0.05$ between different distances and landscape. SE= Standard Error

At Dyamdyam tufts diameters of *T. triandra* and *E. plana* were not significantly different ($P > 0.05$) between the sites. Tuft diameter of *C. dactylon* was slightly highest on near site than middle and far sites. For *S. africanus* tuft diameter was not significant different ($P > 0.05$) across the study sites. At Machibi tuft diameter of *T. triandra* and *C. dactylon* were not significantly differ ($P > 0.05$) between slope and top sites however it was slightly highest on bottom sites. For *S. africanus* tuft diameter was not significantly different ($P > 0.05$) between bottom and slope sites. Tufts diameter of *E. plana* was not significantly different ($P > 0.05$) between all the study sites.

Table 4: Mean tufts diameter (cm) of common grass species found in study areas

Grass species	Dyamdyam				Machibi			
	Near	Middle	Far	SE	Bottom	Slope	Top	SE
<i>T. triandra</i>	2.1 ^a	2.3 ^a	2.0 ^a	0.3	3.0 ^a	2.0 ^b	2.4 ^b	0.4
<i>C. dactylon</i>	3.0 ^a	2.3 ^b	2.4 ^b	0.4	3.4 ^a	2.1 ^b	2.0 ^b	0.7
<i>E. plana</i>	3.0 ^a	3.4 ^a	3.0 ^a	0.3	2.0 ^a	2.0 ^a	2.2 ^a	0.3
<i>S. africanus</i>	3.1 ^a	4.0 ^a	3.0 ^a	0.7	2.2 ^a	1.2 ^b	2.0 ^a	0.5

^{abc}Different superscripts for each species in a row denote significant differences at $p < 0.05$ between different distances and landscape. SE= Standard Error

3.5. Total density of herbaceous species

The results for total mean density of herbaceous species identified in study areas are presented in (Table 5). Both in summer and winter at Dyamdyam total mean density was highest ($P < 0.05$) in far site followed by middle and near. Similarly, at Machibi the total density of herbaceous species was significantly highest in top site followed by slope and bottom in winter and summer. Total density of herbaceous species was highest ($P < 0.05$) in summer than winter across the study sites in both study areas.

Table 5: Mean density (m⁻²) of herbaceous species identified in two study areas.

Areas/Sites	Total Density	Total Density
Dyamdyam	Winter	Summer
Near	30 ^{cb}	43 ^{ca}

Middle	51 ^{Bb}	63 ^{Ba}
Far	67 ^{Ab}	74 ^{Aa}
SE	1.9	1.9
<i>Machibi</i>		
Bottom	33 ^{Cb}	50 ^{Ca}
Slope	57 ^{Bb}	65 ^{Ba}
Top	66 ^{Ab}	72 ^{Aa}
SE	2.2	2.2

^{abc}Lowercase superscripts are used to compare means between seasons within each site. ^{ABC}Uppercase superscripts are used to compare site averages within each season. Means with different superscript within the columns and rows are significantly different ($P < 0.05$).

4. Discussion

4.1. Ecological Status, Life Form and Palatability of Herbaceous Species

In the present study herbaceous species in the study areas were dominated more by increaser and less palatable species. Many studies conducted elsewhere in Africa reported similar findings (Danckwerts, 2001; Sisay & Baars, 2002; Hayes & Holl, 2003; Hein, 2006; Gemedo-Dalle *et al.*, 2006; Morris & Kotze, 2006; Anderson & Hoffman, 2006; Solomon *et al.*, 2007). These findings could be an indication that communal rangelands are overgrazed (Shackleton, 1993) because palatable and decreaser species decline in numbers when the rangeland is over or selectively grazed and replaced by increaser and less palatable species (Sisay & Baars, 2002). The present study also showed that study sites were dominated by perennial species. Similar results were reported by Morris & Kotze (2006). Ruminants can alter vegetation from being dominated by annual species to being dominated by perennials species due to the high stocking rate on communal grazing lands (Smet & Ward, 2003).

4.2. Composition of Common or Dominant Grass Species

Common or dominant grass species identified in the present study were similar to species recorded in the previous study conducted in the Eastern Cape Province of South Africa (Lesoli, 2008), however, they were different in their frequency of occurrence. As observed in the results of this study, it is clear that the proportion of occurrence of highly palatable species (*T. triandra*) is relatively low on near and bottom sites at Dyamyam and Machibi, respectively. These findings suggested that these sites were selectively grazed or overgrazed. This is because palatable grass species are very much under pressure from selective grazing and over utilization and are the first species that are removed under heavily grazed rangelands. This view is supported by Quattrocchi (2006) who reported that, *T. triandra* is an indicator of a good rangeland condition and it quickly disappears in part of rangeland where overgrazing and selective grazing occurs. In addition, *T. triandra* is growing well and most commonly occurs in undisturbed open grasslands with an optimum amount of rainfall. *Cynodon dactylon* was found to be slightly higher on bottom and near sites than other sites. This indicates that bottom and near sites were heavily grazed because *C. dactylon* is the dominant key species in rangeland that is heavily grazed (Van Oudtshoorn, 1992). In addition, *C. dactylon* is the perennial grass that grows well in all types of soil, more especial in sandy soil and disturbed areas (Xu *et al.*, 2011). Frequency of occurrence of *E. plana* and *S. africanus* were relatively low on near and bottom sites compared to other grazing sites. However, these species were expected to be high on these sites because they grow well in disturbed soil such as trampled part of rangeland by livestock as well as near the homestead, roads and drinking areas (Van Oudtshoorn, 2012).

4.3. Biomass Production

The current study showed that mean biomass production in all the study sites was higher in summer than in winter. High biomass production in summer could partly be explained by high rainfall and optimum temperature that promote tropical and subtropical vegetation growth. These results concur with the reports of Noellemeier *et al.* (2006) in semi-arid areas of Argentina and as above Angasa & Oba (2010) in southern Ethiopia, who observed

seasonal variation in biomass production. This was however not in agreement with the study of McDonald *et al.* (1987) who did not find any seasonal variation in biomass production.

Biomass production at Dyamdyam was low on near site from homesteads than the other sites. The present results correspond with earlier findings by Savadogo *et al.* (2007) in the savanna woodlands of Burkina Faso. This can be explained by the fact that near site was heavily grazed due to easy access by ruminants and availability of the drinking areas. Energy requirement of ruminants is different depending on the distance which livestock walk. Therefore, this might lead to declined movement of livestock which in turn would result in high density of livestock on easily accessible grazing sites (Bailey *et al.*, 1996).

In the current study biomass production at Machibi was higher on top and slope sites than bottom sites. Similar results were reported by Senft *et al.* (1985) and Lesoli (2008) in communal grazing lands of South Africa. This indicates that livestock spend most of the time grazing in bottom lands due to easy access. For grazing sites found on heterogeneous landscape, animal grazing distribution pattern might be impacted by variations in landscape (Lesoli, 2008). However, the bottom site in the present study was expected to have high biomass production than slope and top positions due to the high accumulation of nutrients on bottomlands (Coronato & Bertiller, 1996) because nutrients promote plant activities.

4.4. Height and diameter of common or dominant grasses

In the current study all common or dominant grass species, except *S. africanus* at Dyamdyam, had the highest mean height in the middle and far sites rather than the near site. As discussed previously near site had been grazed more intensively than other sites. Therefore, this may cause a short grass height on this site due to high grazing pressure. At Machibi all the common or dominant grass species, except *C. dactylon*, had the highest height on top and slope sites than the bottom site. This is associated with physical damage caused by ruminants. *Cynodon dactylon* had highest height on bottomland than slope. This confirms the fact that *C. dactylon* grow well in overgrazed and disturbed areas (Van Oudtshoorn, 1992) and is common in sandy soil (Xu *et al.*, 2011).

Plant height of *S. africanus*, *T. triandra* and *E. plana* across the study sites and in both study areas was below the maximum growth range of 28–150 cm, 30–150 cm and 40–100 cm respectively suggested by Van Oudtshoorn (2012). Plant height of *C. dactylon* across the study and in both study areas was within the maximum growth range of 5–40 cm reported by Van Oudtshoorn (2012).

This study showed that average tuft diameters of *T. triandra* and *E. plana* at Dyamdyam were not significantly different amongst all the sites. This indicates that a tuft diameter of these species in the present study did not influenced by distance from homestead. However, the difference was expected due to differences in grazing intensity and accumulation of nutrients along the distance gradient from homesteads.

In the current study average tufts diameter of *T. triandra* and *C. dactylon* at Machibi did not show the significant difference between slope and top sites. These results contradict with the report of O' Connor & Pickett (1999) in the semi-arid savannas of East Africa who reported that, there is steady change in grass species along grazing gradients usually characterized by a decline in tuft size. Tufts diameter of *T. triandra* and *C. dactylon* was bigger in bottom site than slope and top sites. This can be attributed to the high accumulation of soil nutrients in the bottomlands (Coronato & Bertiller, 1996) because nutrients promote plant activities. Plant tuft size of *C. dactylon*, *S. africanus*, *T. triandra* and *E. plana* across the study sites and in both study areas was below the maximum tuft diameter range of 5–10 cm, 10–14 cm, 5–15 cm and 5–10 cm respectively reported by (Van Oudtshoorn, 2012).

4.5. Density of herbaceous species

The total mean density of herbaceous species in all the study sites and in both study areas was significantly higher in summer than in winter. Similar results were reported by Bailey *et al.* (1996). This can be attributed by a higher amount of rainfall in summer and the availability of soil nutrients. This is because plant density increases with an increase in rainfall and soil nutrients availability (Ahmad *et al.*, 2007). In the current study, total mean density of herbaceous species strongly influenced by the distance gradient from homesteads. Dyamdyam and Machibi plant density was low on near and bottom sites respectively than other sites in both seasons. Similar results were reported by Senft *et al.* (1985) and Lesoli (2008) in South Africa. High grazing intensity and human activities on bottom and near sites can result in bare patches and veld degradation which in turn results in low

density of herbaceous species (Senft *et al.*, 1985; Lesoli, 2008). Lower plants density on the near and bottom site could also be partially explained by the fact that small stock such as sheep and goats kraaled during night and released in the morning. This could have an effect on plant density during the livestock movement (Lesoli, 2008).

5. Conclusion

The present study showed that most palatable species such as *T. triandra* had low frequency occurrence in near grazing sites than middle and far sites. The study sites were dominated by *T. triandra*, *C. dactylon*, *E. capensis*, *E. plana* and *S. africanus*. The biomass production was significantly higher in summer than in winter season. Density of herbaceous species was low on near grazing site than middle and far. Therefore, it can be concluded that season landscape and distance from homesteads are the most important factors that affect vegetation change and the composition of herbaceous species. Therefore, any veld condition assessment and rangeland management practices in communal rangelands should take into consideration often these factors. In addition, application of suitable grazing systems in communal grazing areas is recommended to inhibit the declining trend of highly palatable species and biomass production near the homestead to improve fodder production for ruminants.

References

- ABULE, E, SNYMAN, H. A. & SMIT, G. N. 2007. Rangeland evaluation in the middle Awash valley of Ethiopia: I. Herbaceous vegetation cover. *Journal of Arid Environment* doi: 10.106/j.jarideny.
- AHMAD, K., HUSSAIN, M., ASHRAF, M., LUQMAN, M., ASHRAF, M. Y. & KHAN. Z. I. 2007. Indigenous vegetation of Soone Valley: At risk of extinction. *Pakistan Journal of Botany* 39 (3): 679–690.
- ANDERSON, P. M. L. & HOFFMAN, M. T. 2006. The impacts of sustained heavily grazing on plant diversity and composition in lowland and upland habitats across the Kamiesberg mountain range in the Succulent Karoo, South Africa. *Journal of Arid Environments*. Doi: 10.106/j.jarideny. 2006–05–17.
- ANGASA, A. & OBA, G. 2010. Effects of grazing pressure, age of enclosures and seasonality on bush cover dynamics and vegetation composition in southern Ethiopia. *Journal of Arid Environments* 74: 111 – 120.
- BAILEY, D. W., GROSS, J. E., LACA, E. A., RITTENHOUSE, L. R., COUGHENOUR, M. B., SWIFT, M. D. & SIMS, P. L. 1996. Mechanisms that result in large herbivore grazing distribution pattern. *Journal of Range Management* 49: 386–400.
- BELSKY, A. J. & BLUEMENTHAL, D. M. 1997. Effects of livestock grazing on Stand Dynamics and Soil in Upland Forest of the Interior West.
- BOND-LAMBERTY, B., WANG, C. & GOWER, S. T. 2002. Net primary production and net ecosystem production of a boreal black spruce wildfire chrosequence. *Global Change Biology* 10: 473–487.
- CORONATO, F. R. & BERTILLER, M. B. 1996. Precipitation and Landscape related effects on soil moisture in semi-arid rangelands of Patagonia. *Journal of Arid Environments* 34: 1–9.
- DANCKWERTS, J. E. 2001. The effects of management practices on livestock performance and veld condition assessment in the semi-arid savannas of Eastern Cape region. *Pasture Research Dohne*. Stutterheim, South Africa [Web:] <http://www.arc.agric.za/institute/rfi/main/projects/project10.htm> [Accessed on 2002/02/02].
- DYKSTERHUIS, E. J. 1994. Condition and management of rangeland based on quantitative ecology. *Journal of Range Management* 2: 104–115.
- EVERSON, T. M. & HATCH, G. P. 1999. Managing veld (rangeland) in communal areas of southern Africa. In: Tainton N M (ed) *Veld Management in South Africa*. University of Natal Press, Pietermaritzburg, pp. 381–388.
- GEMEDO-DALLE, MAASS, B. L. & ISSEETEIN, J. 2006. Rangeland condition trend in the semi-arid Boran lowlands, southern Oromia, Ethiopia. *African Journal of Range and Forage Science* 23 (1): 49–58.
- HAYES, G. E. & HOLL, D. K. 2003. Cattle grazing impacts on annual forbs and vegetation composition of mesic grass veld in California. *Conservation Biology* 17 (6): 1694–1702.
- HEIN, L. 2006. The impact of grazing and rainfall variability on the dynamics of a Sahelian rangeland. *Journal of Arid Environments* 64: 488–504.
- LESOLI, M. S. 2008. Vegetation and soil status, and human perception on the condition of communal rangeland of the Eastern Cape, South Africa. MSc. Thesis, University of Fort Hare, South Africa.
- MAKI, A., KENJI, T., KIYOKAZU, K. & TERUO, H. 2007. Morphological and physico-chemical characteristics of soil in a steppe region of the Kherlen River basin, Mongolia. *Journal of Hydrology* 333: 100–108.
- MCDONALD, L. C., MCFEETERS, R. F., DAESCHEL, M. A. & FLEMING, H. P. 1897. A differential medium for the

- enumeration of homofermentative and heterofermentative lactic acid bacteria. *Applied and Environment Microbiology* 53: 1382–1384.
- MORRIS, C. & KOTZE, D. 2006. Introduction to Veld care (1). Agricultural research council, University of Kwazulu Natal.
- NOELLEMAYER E., QUIROGA, A.R. & ESTELRICH, D. 2006. Soil quality in three range soils of the semi-arid Pampa of Argentina. *Journal of Arid Environments* 65: 142–155.
- O' CONNOR, T. G. & PICKETT, G. A. 1999. The effects of grazing history on the herbaceous composition: population structure and seed banks of some African savanna grasslands. *Journal of Applied Ecology* 29 (1):247–260.
- O' FARRELL, P. L., DONALDSON, J. S. & HAFFMAN, M. T. 2007. The influence of ecosystem goods and services on livestock management practices on the Bokkeveld plateau, South Africa. *Agriculture, ecosystem and environment* doi: 10.1016/j. agee. 2007-.01–025.
- OZTAS, T., KOC, A. & COMAKLI, B. 2003. Changes in vegetation and soil properties along a slope on overgrazing and eroded rangelands. *Journal of Arid Environments* 55: 93–100.
- QUATTROCCHI, U. 2006. CRC World dictionary of grasses: common name, scientific names, eponyms, synonyms and etymology. CRC Press, Taylor and Francis Group, Boca Raton, USA.
- SAVADOGO, P., SAWADOGO, L. & TIVEAU, D. 2007. Effects of grazing intensity and prescribed fire on soil physical and hydrological properties and pasture yield in the savanna woodlands of Burkina Faso. *Agriculture, Ecosystems and Environment* 11: 80 – 92.
- SENF, R. L., RITTENHOUSE, L. R. & WOODMANSEE, R. G. 1985. Factors influencing patterns of cattle grazing behaviour on short grass steppe. *Journal of Range management* 38: 82–87.
- SHACKLETON, C. M. 1993. Fuelwood harvesting and sustainable utilization in a communal grazing land and protected area of eastern Transvaal Lowveld. *Biological Conservation* 63: 247– 257.
- SISAY, A. & BAARS, R. M. T. 2002. Grass composition and rangeland condition of the major grazing areas in the mid Rift Valley, Ethiopia. *African of range and forage Science* 19: 161–166.
- SMET, M. & WARD, D. 2005. A comparison of the effects of different rangeland management system on plant species composition, diversity and vegetation structure in a Semi-Arid Savanna. *African Journal of Range and Forage Science* 22(1): 59–71.
- SOLOMON, T. B., SNYMAN, H. A. & SMIT, G. N. 2007. Cattle-rangeland management practices and perceptions of pastoralists towards rangelands degradation in the Boran zone of Southern Ethiopia. *Journal of Environmental Management* 82: 481–494.
- TROLLOPE, W. S. W., TROLLOPE, L. A. & BOSCH, O. J. H. 1990. Veld and pasture management terminology in Southern Africa. *Journal of Grassland Society of Southern Africa* 7: -52–61.
- VAN OUDTSHOORN, F. V. 1992. Guide to Grasses of South Africa. Pretoria: Briza Publications. First edition
- VAN OUDTSHOORN, F. V. 2012. Guide to Grasses of southern Africa. Briza Publications. Third revised edition.
- VORSTER, M. 1982. The development of the ecological index method for assessing veld condition in the Karoo. *Proceeding of the Grassland Society of Southern Africa* 17: 84–89.
- XU, J., WANG, Z. & CHENG, J.J. 2011. Bermuda grass as feedstock for biofuel production: A review. *Bio-resource Technology* 102: 7613–7620.